
THE AIR TOXICS PROGRAM

What are air toxics?

- Pollutants which...
 - may cause cancer or other serious health effects in humans or in the environment from both short and long term exposure
 - may disperse locally, regionally, nationally, or globally
 - after deposition may persist in the environment and/or bioaccumulate in the food chair
 - possess a variety of physical and chemical characteristics that enhance potential for multi-media exposure
- > 188 compounds listed in the Clean Air Act




What health & environmental effects do they cause?

- More than half are known or suspected to be human carcinogens
- Many known to affect respiratory, neurologic, immune, hormonal, or reproductive systems
- More susceptible or sensitive populations at greater risk
- Known to have similar effects in many fish and animal species
- Environmental effects may be felt by individual species within ecosystem or by entire ecosystem where affected species are found

Source of air toxics

- Literally, thousands of sources
- Stationary sources include large industrial complexes
 - chemical plants, oil refineries, and steel mills
- Area sources which are small stationary sources
 - dry cleaners, gas stations, and small manufacturers
- Mobile sources
 - on-road vehicles cars, trucks, and buses
 - non-road vehicles ships, lawn mowers, and farm equipment




History of the Program

1970 CAA Section 112 Provisions

Directed EPA to identify air pollutants with hazardous effects and establish standards to prevent any adverse effects "with an ample margin of safety"

Prior to 1990, EPA set standards for only 8 pollutants:

arsenic	beryllium	radon
asbestos	mercury	radionuclides
benzene	vinyl chloride	




Section 112: Statutory Provisions

- Sec 112(a) Definitions
- Sec 112(b) List of Pollutants
- Sec 112(c) List of Source Categories
- Sec 112(d) Emission Standards
- > Sec 112(e) Schedule for Standards & Review
- Sec 112(f) Protect Health & Enviro Standard
- Sec 112(g) Modifications
- Sec 112(h) Work Practice Standards
- Sec 112(i) Schedule for Compliance
- Sec 112(j) Equiv. Emission Limitation by Permit

Section 112: Statutory Provisions (continued)

- Sec 112(k) Area Source Program
- Sec 112(I) State Programs
- Sec 112(m) Atmospheric Deposition to Great Lakes & Coastal Waters
- Sec 112(n) Other Provisions
- Sec 112(o) National Academy of Sciences Study
- Sec 112(p) Mickey Leland Urban Air Toxics Rsh Ctr
- Sec 112(q) Savings Provision
- Sec 112(r) Prevention of Accidental Releases
- Sec 112(s) Periodic Report




Air Toxics Provisions

- Statutory list of 189 HAPs
 - EPA can delete or add to list
- List of categories of industrial facilities emitting substantial quantities of HAPs
- Establish MACT for each category
- Expected significant reductions below existing emission limits
- All standards must be promulgated by November 15, 2000


Source Category List

- Section 112(c) requires list of major sources emitting:
 - >10 tons/year of one pollutant
 - >25 tons/year of a combination of pollutants
- > EPA published initial list July 16, 1992
 - 166 categories of major sources
 - 8 categories of area sources
- List can be revised
- Urban strategy will add more area sources




National Standards

- Section 112(d)
 - For new sources, best controlled similar source
 - For existing sources, the average emission limitation achieved by the best performing 12%
 - Additional reductions based on costs and non-air quality impacts



Case-by-Case MACT


- Section 112(j) the "permit hammer" provisions
 - Occurs if MACT is not promulgated within 18 months of statutory deadline
 - Permit contains case-by-case MACT
- Section 112(g) sources with increases in HAP emissions
 - Changed operations
 - Offset and de minimis allowances



Goals of the Program


Overall Goal

Reduce risks to the population and minimize environmental impacts from air toxics




Current GPRA Goal

➤ By 2010, reduce air toxics emissions by 75% from 1993 levels to significantly reduce the risks of the population to cancer and other serious adverse health effects caused by airborne toxics



Future GPRA Goal (draft)


▶ By 2015-2020, eliminate unacceptable risks of cancer and other significant health problems from air toxics emissions for at least 95% of the population and substantially reduce or eliminate adverse effects on our natural environment



Status of the Air Toxics Program

The Air Toxics Program

- Designed to characterize, prioritize, and equitably address the serious impacts of hazardous air pollutants on public health and the environment through a strategic combination of:
 - regulatory approaches
 - voluntary partnerships
 - ongoing research and assessments
 - education and outreach



Components of the Air Toxics Program

- Source-specific standards and sector-based standards
 - MACT (Sections 112 and 129)
 - Residual Risk
 - Utilities Study
- National, regional, community-based initiatives to focus on multimedia and cumulative risks
 - Integrated Urban Air Toxics Strategy
 PBT and TMDL initiatives
 - Great Waters
 - Mercury initiatives
- National air toxics assessments (NATA)
 - Emission inventories
- Air quality, exposure, and risk modeling
- Monitoring network
- Ongoing research on effects and assessment

Clean Air Partnerships

> Educational outreach tools

The MACT Program

MACT: Section 112 Program

- 188 hazardous air pollutants initially listed in CAAA
- 174 source categories listed by EPA

Source categories divided into bins:


BIN	STATUTORY DATE	STANDARDS/ SOURCE CATEGORIES	PROPOSED TO DATE	COMPLETED TO DATE
2 year	11/15/92	2 standards 6 source cat.		2 standards 6 source cat.
4 year	11/15/94	18 standards 40 source cat.		19 standards 40 source cat.
7 year	11/15/97	30 standards 41 source cat.		24 standards 35 source cat. 5 delisted
10 year	11/15/00	62 standards 94 source cat.	4 standards 4 source cat.	1 standard 1 source cat. 2 delisted

MACT: Status of Section 112

- Completed all 2 and 4 year standards
 - 21 standards covering 46 source categories
- Completed all 7 year standards
 - 24 standards covering 35 source categories
 - Delisted 5 source categories
- Initiated work on all 10 year standards
 - Total of 62 standards covering 94 source categories
 - Promulgated 1 standard covering 1 source category
 - Proposed 4 standards covering 4 source categories
 - Delisted 2 source categories

MACT: Section 129 Combustion Standards


- Municipal Waste Combustion
 - Initially promulgated December 1995
- Hospital/Medical/Infectious Waste Incineration
 - Promulgated September 1997
- Promulgate by November 2000:
 - Industrial/Commercial Waste Incinerators
 - Small MWC Units



MACT: Implementation Status

► Regions have delegated 69% of the major source standards and 49% of area source standards


► The "apparent" national aggregate MACT compliance rate is 36%



The Residual Risk Program


Residual Risk: Purpose

- Assess post-MACT risks from source categories
- Set additional standards, if needed, to protect public health with an "ample margin of safety"
 - within 8 years of MACT promulgation
- Prevention of "adverse ecological impact" with consideration of other factors
- Report to Congress March1999



Residual Risk: Need for a Risk Management Framework

- Source categories will likely require unique analyses
 - Data availability and quality
 - Various types of source categories
 - number of sources
 - HAPs emitted
- Enable consistent decision process




Residual Risk Management Framework

- Will Be
 - A broad conceptual framework which identifies:
 - decision points in the process
 - major inputs into these decisions
 - type of information required to support each decision
 - guidance for decision making under uncertainty
- > Will MOT be
 - A prescriptive decision tool outlining rigid criteria for each decision


Residual Risk: Risk Management Decisions

- Protection of public health with "ample margin of safety"
 - Cancer (Benzene NESHAP Framework)
 - Noncancer
- Prevention of "adverse environmental impact" with consideration of other factors




Residual Risk: Benzene NESHAP Ample Margin of Safety (AMS) Framework

- CAA maintained the use of AMS as applied in Benzene NESHAP
- > Two-step process
 - safe or acceptable
 - ample margin of safety
- Currently defined for Benzene NESHAP (linear carcinogen)
- Must apply principles to others as well
 - non-linear carcinogen
 - noncarcinogens



Residual Risk: Overview of Benzene NESHAP Relevant Risk Ranges



Residual Risk: General Framework for Assessment Process

- Use a tiered approach to conducting assessments:
 - Data Collection & Problem Formulation
 - initial scoping plus data collection and re-scoping, as necessary, prior to each level of analysis
 - Screening Analysis
 - conservative, relying on readily available information and defaults
 - More Refined Analyses
 - Increasingly more realistic and more resource intensive
 - Utilize different tools or data

Residual Risk: 2 & 4 Year MACT Standards


- Due 2001
 - Coke Ovens*
- Due 2002
 - Dry Cleaning*
 - EO Sterilizers*
 - Gas Distribution*
 - Halogenated Solvent Cleaning*
 - Industrial Cooling Towers*
 - Magnetic Tape*

- Due 2003
 - Aerospace Manuf.
 - Chrome Electroplating*
 - Petro Refineries*
 - Polymers and Resins II*
 - Sec. Lead Smelters*
 - Shipbuilding
 - Wood Furniture
 - Marine Vessel Loading*
 - HON*
- Due 2004

 - Printing/PublishingPolymer & Resins I, I\
 - Offsite Waste


Residual Risk Assessments -- Status

- > 12 Source Categories have been started
 - Screening assessments underway on 8
 - More refined assessments underway on 4




Residual Risk Assessments: Status (continued)

- Characteristics of source categories vary
 - number of sources range from 23 to >5,000
 - majority have fewer than 100 sources
 - number of HAPs range from 1 to over 100
 - 3 have > 50 HAPs
 - Carcinogenic and noncarcinogenic HAPs emitted
 - Some potential for multipathway exposures




Residual Risk: Status -- Toxicity Data

- Hierarchy of data used for developing dose-response data for risk assessment
 - IRIS
 - ATSDR
 - California EPA
 - HEAST
 - Other governmental agencies
- ► As a result, we have toxicity data for 150 of the 188 HAPs




Residual Risk Assessments: Status -- Data

- Data used
 - MACT Background Information Document, NTI
 - State and Regional Offices
 - Industry
- Requests for data from industry made for 6 source categories
 - data have been provided from industry on 2 source categories
- In the absence of data:
 - use conservative assumptions
 - use Section 114 authorities



Residual Risk: Status -- Tools


- Range of tools used in risk assessments
 - SCREEN 3 modeling conducted for 3 source categories
 - ISC/HEM applied to 6 source categories, to date
 - IEM applied to 1 source categories date, 4 more anticipated
 - TRIM being developed and will be available on a limited basis in the next year



Great Waters Program


Great Waters Program

- Section 112(m) Requirements
 - Deposition assessment
 - Periodic Report to Congress
 - Regulatory determination
 - Monitoring for Great Lakes, Chesapeake Bay, and Coastal Waters
- > 15 Pollutants of Concern
 - Mercury, lead, and cadmium
 - Nitrogen compounds
 - POM/PAHs, dioxins, and furans
 - PCBs and 7 banned/restricted pesticides




Great Waters Program: Accomplishments

- Two Reports to Congress
 - Issued May 1994 and June 1997
- Adequacy determination supporting Section 112 authorities
 - Issued March 1998
- Third Report to be issued summer 2000



Great Waters Program: Current Directions

- Primary focus on nitrogen and mercury
- Developing monitoring capabilities of NEPs
 - workshops and special studies
- Developing analytical tools and legal strategies
- Building partnerships




Great Waters Program: Current Projects

- Great Lakes monitoring/studies
- Hg Deposition and TMDL Pilot
 - Florida Everglades and Devils Lake, WI
- Monitoring Network Long & Short Term
- Air Emissions from Concentrated Animal Feeding Operations (CAFOs)
- Develop better communication tools
 - website for better data access
 - handbook for air dep studies
- Develop Action Plan
 - complete action plan in 2000 and begin implementation in 2001

Utility Study

Utility Study

- Study completed in February 1998
- Currently collecting information to support regulatory determination
 - issued information collection request (ICR) to utility industry
 - conducting (through ORD) analysis of potential control technologies
 - continue analysis of health-related issues
- Regulatory determination by 12/15/00




Urban Air Toxics Strategy

Urban Air Toxics Strategy


- Final Strategy Issued July 19, 1999
 - list of 33 HAPs
 - list of 13 new area source categories
 - 202(I) standards and low sulfur diesel regulations
 - activities for residual risk and 10 year MACT
- Initiate mobile source standards first
 - propose July 2000
 - promulgate December 2000

Area source standards by 2004



Urban Air Toxics Strategy: Goals

- > 75% reduction in cancer "incidence"
 - Scope: national, from all stationary sources looking at all HAP
- "Substantial" reduction in noncancer "risks"
 - Scope: national, only from area sources, looking at all noncarcinogenic HAP
- Address disproportionate risks




Urban Air Toxics Strategy: Progress Assessment

- Iterative Analyses
 - Preliminary -- comparison of 1990 and 1996 emissions and ambient monitoring data
 - Periodic assessments -- initially based on 1996 emission inventory
- Allow for evolution of methods over time
 - Improved tools and data
 - Progression from screening level to more complex assessment


Urban Air Toxics Strategy: Area Source Standards

- Options: How to develop standards
 - MACT
 - Traditional GACT
 - Flexible GACT
- Options: Where standards apply
 - National standards apply everywhere
 - National standards apply only in urban areas
 - State and local standards



Urban Air Toxics Strategy: Other Activities


- Identify research needs
- Work in partnerships with State, local and tribal governments to develop programs that address goals of urban strategy
- Federal Advisory Committee Act (FACA)
- Initiate pilot projects with the Office of Indoor Air
- Education and outreach



Indoor Air


Indoor Air

- Both indoor and outdoor air exposures to air toxics are important
- Indoor air and outdoor air are interdependent
 - outdoor air toxics set baseline level indoors
 - outdoor air toxics infiltrate or are ventilated indoors
 - indoor emissions are ventilated outdoors



Indoor Air: Integrated Strategy


- ► For meaningful reduction of risks, we must address **BOTH** indoor and outdoor sources of air toxics
- > Will do this through:
 - timing of joint strategies to reduce air toxics
 - joint communications and coordination
 - joint project(s)
 - pilot study in one community



Mobile Source Strategy


Mobile Source Strategy

- Mobile sources are significant contributors to urban toxics emissions inventories
- Many compounds and elements are known to be emitted by motor vehicles
- ► There are 3 primary ways to reduce emissions of mobile source air toxics:
 - fuel controls
 - engine controls
 - reduction of vehicle miles traveled




Mobile Source Strategy: Standards Under Development

- Heavy Duty Diesel Engines (mid 2000)
 - final rule that reconfirms 2004 standards for heavy duty diesels
 - proposal to control sulfur in diesel fule fuel and establish more stringent standards for heavy duty vehicles and engines
- Tier 2 Rule (finalized December 1999)
 - new stringent emission standards and gasoline sulfur controls from light duty vehicles and trucks



Mobile Source Strategy: Standards Under Development (continued)


- Tier 3 Non-Road Diesel
 - propose late 2000/finalize 2001
 - non-road diesel fuel controls and emission standards for non-road diesel engines
- Section 202(I) Rule
 - propose July 2000/finalize December 2000
 - will designate motor vehicle air toxics and consider control options




National Air Toxics Assessment (NATA)

NATA: What is it?

- Development of analytical tools to:
 - identify areas of concern
 - improve characterization of risks and risk reductions for stationary and mobile sources
 - track our progress and prioritize efforts
- To be accomplished through:
 - emission inventories development
 - monitoring networks
 - modeling




NATA 1996 National Air Toxics Screening Assessment




NATA Screening Assessment: What it is...

- Identification of priority hazardous air pollutants and sources
- Identification of potential national scale air toxics problems
- Prioritization of future data collection and localized modeling efforts
- Allow for better analyses
- Measure progress against national goals
- > Help set research agenda




NATA Screening Assessment: What it is NOT...

- > NOT called CEP2
- ► NOT the only thing included under the NATA umbrella
- NOT being used to make specific regulatory decisions




NATA Screening Assessment: Scope and Schedule

- National Effort: Continental US, Virgin Islands, and Puerto Rico
- Addresses emissions for 33 priority HAPs plus diesel PM
- > Phased roll-out
 - 1st Air Quality
 - 2nd Exposure Models
 - 3rd Risk Analysis




NATA Screening Assessment: Scope and Schedule *(continued)*

- Provide initial result summaries (emissions and ambient estimates) and model-to-monitor comparisons on website spring 2000
- Provide draft final results (exposures and risks) on website summer 2000 for public and peer review
- > Final peer reviewed assessment by December 2000




NATA: Public Health Implications

- Scope
 - 33 Urban HAPs
 - County resolution
 - Inhalation only (not multipathway)
 - Chronic effects only (not acute)
 - Characterization will address these issues qualitatively
- Purpose
 - Screening-level assessment




NATA: Risk Characterization Outputs

- > WILL produce:
 - County-level cancer risk and noncancer hazard for each HAP
 - County-level cumulative risk and hazard for each HAP
- ► WILL be clearly identified as "screening-level" for use in:
 - Setting the research agenda
 - Measuring progress against national goals
 - Identifying priority HAP & sources



NATA: Risk Characterization Outputs *(continued)*


- > WILL NOT produce:
 - Reliable risk estimates below county level
 - Information on hot spots
- ► The screening-level characterization should **NOT** be used for:
 - Comparing risks at the local scale
 - Regulatory decision making
 - Showings of harm or absence of harm



Partnerships

Develop Partnerships

- > State/Locals/Tribes
- > Other EPA Offices
- > Other Health Agencies
 - OSHA, CDC, NIEHS
- Industry
- Public Interest Groups



For More Information on Air Toxics

- Visit the following websites:
 - EPA's Office of Air and Radiation (OAR)
 - www.epa.gov/oar
 - Unified Air Toxics Website (UATW)
 - www.epa.gov.ttn/uatw
 - OAR Policy & Guidance Website
 - provides access to OAR rules, policies, and guidance documents
 - www.epa.gov/ttn/oarpg

In Summary

- Continue to implement air toxics program to meet GPRA goals...
 - complete 10-year MACT standards
 - effectively implement and enforce standards
 - continue residual risk analysis of promulgated rules
 - develop/implement urban based strategies
 - develop data and tools to enhance our capabilities to characterize risk and monitor progress
 - look at multi-media impacts
 - partnership fund
 - mobile sources strategies/standards
 - indoor air strategies

